

Michigan
Adult Treatment Protocols
ANAPHYLAXIS/ALLERGIC REACTION

Date: May 31, 2012

Page 1 of 2

Anaphylaxis/Allergic Reaction

Pre-Medical Control

MFR/EMT/SPECIALIST/PARAMEDIC

1. Follow **General Pre-hospital Care Protocol**.
2. Determine substance or source of exposure, remove patient from source if known and able.
3. Assist the patient in administration of their own epinephrine auto-injector, if available.

EMT/SPECIALIST

4. In cases of severe allergic reaction, wheezing or hypotension, administer epinephrine via auto-injector.
5. Albuterol may be indicated. Refer to **Nebulized Bronchodilators Procedure**.

SPECIALIST/PARAMEDIC

6. Administer a NS IV/IO fluid bolus up to 1 liter, wide open as indicated.

PARAMEDIC

7. If patient is symptomatic, administer diphenhydramine 50 mg IM or IV/IO.
8. In cases of severe allergic reaction, wheezing or hypotension:
 - A. Administer Epinephrine 1:1000, 0.3 mg (0.3 ml) IM OR via auto-injector.
9. In cases of profound anaphylactic shock (near cardiac arrest):
 - A. Administer Epinephrine 1:10,000, 0.3 mg (3 ml) slow IV/IO.
10. Per MCA selection, administer Bronchodilator per **Nebulized Bronchodilators Procedure**.
11. Per MCA Selection, administer Prednisone **OR** Methylprednisolone.

<u>Medication Options:</u>	
<u>Prednisone</u>	
50 mg tablet PO	
<input checked="" type="checkbox"/> YES	<input type="checkbox"/> NO
<u>Methylprednisolone</u>	
125 mg IV	
<input checked="" type="checkbox"/> YES	<input type="checkbox"/> NO

Post-Medical Control:

EMT/SPECIALIST

1. Additional Epinephrine via auto-injector.

PARAMEDIC

2. Additional Epinephrine 1:1,000, 0.3 mg (0.3 ml) IM; or Epinephrine 1:10,000 0.3 mg (3ml) slow IV/IO if critically ill (near cardiac arrest).

Michigan
Adult Treatment Protocols
ANAPHYLAXIS/ALLERGIC REACTION

Date: May 31, 2012

Page 2 of 2

