

Michigan
Adult Cardiac Protocols
ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA)

Date: May 31, 2012

Page 1 of 2

Asystole / Pulseless Electrical Activity

During CPR, consider reversible causes of Asystole/PEA and treat as indicated. Causes and efforts to correct them include but are not limited to:

- Hypovolemia – give NS IV/IO fluid bolus up to 1 liter, wide open.
- Hypoxia – reassess airway and ventilate with high flow oxygen
- Tension pneumothorax – see **Pleural Decompression Procedure**
- Hypothermia – follow **Hypothermia Cardiac Arrest Protocol** rapid transport
- Hyperkalemia (history of renal failure) – see #5 below.

Pre-Medical Control
PARAMEDIC

1. Follow the **Cardiac Arrest - General Protocol**.
2. Confirm that patient is in asystole by evaluating more than one lead.
3. Administer Epinephrine 1:10,000, 1 mg (10 ml) IV/IO, repeat every 3-5 minutes.
4. Per MCA selection, administer Vasopressin 40 Units IV/IO in place of second dose of Epinephrine.

<u>Vasopressin</u> 40 Units IV/IO
<input type="checkbox"/> Included
<input checked="" type="checkbox"/> Not Included

5. In a dialysis patient hyperkalemia is likely. Administer Calcium Chloride 1gm IV/IO and Sodium Bicarbonate 1 mEq/kg IV/IO with 20 ml NS flush in between medications.
6. Continue CPR and reassess rhythm every 2 minutes.

Michigan
Adult Cardiac Protocols

ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA)

Date: May 31, 2012

Page 2 of 2

During CPR, consider reversible causes of Asystole/PEA and treat as indicated. Causes and efforts to correct them include but are not limited to:

- Hypovolemia – give NS IV/IO fluid bolus up to 1 liter, wide open.
- Hypoxia – reassess airway and ventilate with high flow oxygen
- Tension pneumothorax – See **Pleural Decompression Procedure**
- Hypothermia – follow **Hypothermia Cardiac Arrest Protocol** rapid transport
- Hyperkalemia (history of renal failure) – see below

- Follow **Cardiac Arrest – General Protocol.**
- Confirm that patient is in asystole by evaluating more than one lead.

Vasopressin

40 Units IV/IO

Included

Not Included

- Administer Epinephrine 1:10,000, 1 mg (10 ml) IV/IO, repeat every 3-5 minutes.
- Per MCA selection, administer Vasopressin 40 units IV/IO in place of second dose of Epinephrine

- In a dialysis patient hyperkalemia is likely. Administer Calcium Chloride 1 gm IV/IO and Sodium Bicarbonate 1 mEq/kg IV/IO with 20 ml NS flush in between medications.
- Continue CPR, reassess rhythm every 2 minutes.

**Contact
Medical
Control**